

Constrained Reductions of 2D dispersionless Toda Hierarchy, Hamiltonian Structure and Interface Dynamics.

J. Harnad¹, I. Loutsenko², O.Yermolayeva³

¹CRM, Universite de Montreal, C.P. 6128, succ. centre-ville, Montreal, Quebec, H3C 3J7, Canada.
e-mail: harnad@crm.umonreal.ca

²Institute of Mathematics, University of Oxford, 24 - 29 St. Gilles', Oxford, OX1 3LB, England
e-mail: loutsenk@maths.ox.ac.uk

³SISSA, Via Beirut 2-4, 34014, Trieste, Italy, e-mail: yermola@fm.sissa.it

Abstract

Finite-dimensional reductions of the 2D dispersionless Toda hierarchy constrained by the “string equation” are studied. These include solutions determined by polynomial, rational or logarithmic functions, which are of interest in relation to the “Laplacian growth” or Hele-Shaw problem governing interface dynamics. The consistency of such reductions is proved, and the Hamiltonian structure of the reduced dynamics is derived. The Poisson structure of the rationally reduced dispersionless Toda hierarchies is also derived.

1 Introduction.

This paper concerns rational and logarithmic reductions of the 2D dispersionless Toda hierarchy of integrable equations (henceforth 2dToda). The subject is motivated by important applications to problems in interface dynamics and statistical physics.

Laplacian growth is a process that governs the dynamics of the boundary in the plane separating two disjoint, open regions $\mathcal{D}_+$ and $\mathcal{D}_-$ in which harmonic (scalar) fields are defined. These may be interpreted as the pressure fields for two incompressible viscous fluids (Hele-Shaw problem). The movement of the boundary is determined (according to Darcy’s law, in the case of viscous fluids) by equating the normal velocity of the boundary to the boundary value of the gradient of the field. In particular, one region (say, the “interior” region $\mathcal{D}_+$) may be chosen to be bounded and have constant harmonic field (corresponding to zero viscosity) with the boundary condition for the “exterior” $\mathcal{D}_-$ region at infinity to be such that there is a unit sink, implying that the area of the interior region grows linearly in time [16]. Denoting the harmonic field (e.g. the pressure) in the exterior region by $P(X, Y)$, this satisfies the conditions:

$$\begin{aligned} \Delta P(X, Y) &= 0 \\ P &\rightarrow (4\pi)^{-1} \ln(X^2 + Y^2) \quad \text{as} \quad X^2 + Y^2 \rightarrow \infty \end{aligned} \quad (1)$$

in the cartesian coordinates (X, Y) . The normalized exterior normal velocity at the boundary is given by:

$$v_n = -n \nabla P. \quad (2)$$

where n denotes an outward normal to $\partial\mathcal{D}_+$. In the case where the boundary is an analytic curve it is usual to use the Riemann mapping theorem to introduce a time-dependent conformal map from the exterior of the unit circle in the complex w plane to the exterior region $\mathcal{D}_-$ in the “physical” plane $z = X + iY$ taking the unit circle to the boundary $\partial\mathcal{D}_+$.

$$z = z(w, x), \quad w = \exp(i\phi), \quad 0 < \phi < 2\pi, \quad (3)$$

where x stands for the physical time of the Hele-Shaw problem. We choose this unusual notation for time for consistency with that used in the literature on the dispersionless integrable systems.

Simple considerations [15], [4], [12] show (see Appendix A) that equations (1, 2) are equivalent to the following equation

$$\text{Im} \left(\frac{\partial z}{\partial \phi} \frac{\partial \bar{z}}{\partial x} \right) = w \left(\frac{\partial z(w, x)}{\partial w} \frac{\partial \bar{z}(1/w, x)}{\partial x} - \frac{\partial z(w, x)}{\partial x} \frac{\partial \bar{z}(1/w, x)}{\partial w} \right) = 1, \quad (4)$$

where bar stands for complex conjugation (and $\bar{w} = w^{-1}$ on the boundary curve). In our notations $\bar{z}(w) = \sum_i \bar{z}_i w^i$ if $z(w) = \sum_i z_i w^i$, while $\bar{z}(w) = \sum_i \bar{z}_i \bar{w}^i$.

Known as the Galin-Polubarinova equation in the Hele-Shaw problem, eq (4) plays an essential role in the theory of infinite-dimensional integrable hierarchies in the dispersionless limit. The relation between the boundary dynamics above and the dispersionless limit of the integrable Toda hierarchy constrained by (4) was shown in [12].

Equation (4) may be interpreted as a constraint on an infinite commuting set of dynamical systems defined in the space $z(x, w)$ of one-parameter families of conformal maps. This constraint represents fixed points of an “additional symmetry” [13] and is called the “string equation” in the theory of integrable systems. The most interesting aspect of such constrained 2dToda flows is that they admit finite-dimensional reductions, which include so-called “multi-finger” solutions [7]. These solutions are of great importance in practical applications and describe numerous phenomena, such as viscous fingering in a Hele-Shaw cell [7],[10],[17] and pattern formation in the quantum Hall effect [1].

In what follows, we consider finite-dimensional reductions of (4) in the context of the 2dToda hierarchy. We first study algebraic solutions of the problem, ignoring the real structure and treating z , $\bar{z}$ as independent functions, and w as a formal variable. Returning to the applications to interface dynamics we identify bar with complex conjugation.

2 2dToda hierarchy and string equation.

The 2dToda hierarchy is a dispersionless limit [18] of the two-dimensional Toda hierarchy and is defined in terms of two functions $z(w, x)$ and $\bar{z}(w^{-1}, x)$ of the form:

$$z(w, x) = r(x)w + \sum_{k=0}^{\infty} u_k(x)w^{-k}, \quad (5)$$

$$\bar{z}(w^{-1}, x) = r(x)w^{-1} + \sum_{k=0}^{\infty} \bar{u}_k(x)w^k. \quad (6)$$

The 2dToda flow equations are

$$\begin{aligned} \partial_{t_k} z &= \{H_k, z\} & \partial_{\bar{t}_k} \bar{z} &= \{\bar{H}_k, \bar{z}\} \\ \partial_{t, k} \bar{z} &= \{H_k, \bar{z}\} & \partial_{\bar{t}_k} z &= \{\bar{H}_k, z\} \end{aligned} \quad (7)$$

where the “Poisson-Lax” bracket notation here denotes

$$\{f, g\} := w \frac{\partial f}{\partial w} \frac{\partial g}{\partial x} - w \frac{\partial f}{\partial x} \frac{\partial g}{\partial w} \quad (8)$$

and the coefficients $r(x), u_k(x), \bar{u}_k(x)$ are viewed as coordinate functions on the phase space. The evolution functions are defined as follows

$$H_k = (z^k)_+ + 1/2(z^k)_0, \quad \bar{H}_k = (\bar{z}^k)_- + 1/2(\bar{z}^k)_0, \quad (9)$$

where subscripts $\pm, 0$ denote the negative/positive and zero parts of the formal Laurent expansion in w (i.e. $f_+ = \sum_{i>0} f_i w^i$, $f_- = \sum_{i<0} f_i w^i$ if $f = \sum_i f_i w^i$).

It is important to note that, despite the notation, (8) is not really a Poisson bracket defined on the infinite-dimensional phase space of the 2dToda system (7) (with coordinates $r, u_k, \bar{u}_k$), but rather a “quasiclassical” $\hbar \rightarrow 0$ limit of commutators in the Lax representation of the dispersive version of (7)

(where $\hbar$ stands for the lattice spacing, see [18]). Values (9) are dispersionless analogs of the upper or lower diagonal parts of the powers of Lax matrices.

The 2d Toda system is nevertheless an integrable Hamiltonian system of PDEs for the functions $r(x, t_1, \dots, \bar{t}_1 \dots)$, $u_k(x, t_1, \dots, \bar{t}_1 \dots)$, $\bar{u}_k(x, t_1, \dots, \bar{t}_1 \dots)$, obtained by equating coefficients of (7) as Laurent polynomials in the dummy variable w . One can easily verify that vector fields (7) commute, i.e

$$\frac{\partial^2}{\partial t_i \partial t_j} \begin{pmatrix} z \\ \bar{z} \end{pmatrix} = \frac{\partial^2}{\partial t_j \partial t_i} \begin{pmatrix} z \\ \bar{z} \end{pmatrix}, \quad \frac{\partial^2}{\partial \bar{t}_i \partial t_j} \begin{pmatrix} z \\ \bar{z} \end{pmatrix} = \frac{\partial^2}{\partial t_j \partial \bar{t}_i} \begin{pmatrix} z \\ \bar{z} \end{pmatrix}, \quad \frac{\partial^2}{\partial \bar{t}_j \partial \bar{t}_i} \begin{pmatrix} z \\ \bar{z} \end{pmatrix} = \frac{\partial^2}{\partial \bar{t}_i \partial \bar{t}_j} \begin{pmatrix} z \\ \bar{z} \end{pmatrix}$$

due to the "zero curvature" conditions

$$\frac{\partial H_i}{\partial t_j} - \frac{\partial H_j}{\partial t_i} - \{H_i, H_j\} = 0, \quad \frac{\partial H_i}{\partial \bar{t}_j} - \frac{\partial \bar{H}_j}{\partial \bar{t}_i} - \{H_i, \bar{H}_j\} = 0, \quad \frac{\partial \bar{H}_i}{\partial \bar{t}_j} - \frac{\partial \bar{H}_j}{\partial \bar{t}_i} - \{\bar{H}_i, \bar{H}_j\} = 0,$$

which follows from equations of motion (7) and definitions (8),(9).

The Galin-Polubarinova equation (4) for the Hele-Shaw problem written in terms of the Poisson-Lax bracket (8)

$$\{z(w, x), \bar{z}(w^{-1}, x)\} = 1 \quad (10)$$

is the string equation.

It is fundamental that equation (10) is invariant under the flows generated by (7). Indeed

$$\partial_{t_k} \{z, \bar{z}\} = \{\partial_{t_k} z, \bar{z}\} + \{z, \partial_{t_k} \bar{z}\} = \{\{H_k, z\}, \bar{z}\} + \{z, \{H_k, \bar{z}\}\} = \{\{\bar{z}, z\}, H_k\} = 0, \quad (11)$$

where we have used (7), (10) and the Jacobi identity for the Lax-Poisson bracket (8).

Thus the string equation (10) defines an invariant manifold under the Toda flows (7) and a reduction of the Toda hierarchy. On the other hand, the Toda flows may be viewed as symmetries generating new solutions of Eq (10).

3 Reductions of the 2d Toda hierarchy constrained by the string equation.

The reduction of the 2d Toda hierarchy by the string equation is still a compatible set of infinite-dimensional dynamical systems.

Indeed, as seen from (5)-(6) the string equation (10) is a system of ODE's of the form

$$dr/dx = R(r, u_1, \dots, \bar{u}_1, \dots), \quad du_i/dx = U_i(r, u_1, \dots, \bar{u}_1, \dots), \quad d\bar{u}_i/dx = \bar{U}_i(r, u_1, \dots, \bar{u}_1, \dots) \quad (12)$$

In what follows we will be interested in further "functional" reductions where $z, \bar{z}$ are polynomial, rational or logarithmic functions of w . As shown below, such reductions are consistent with (7) (i.e. the corresponding ansatz for z is preserved by the 2D Toda flows) if the string equation (10) holds. Thus, for consistency we need a double ("functional" plus "string") reduction. These pairs of reductions define finite-dimensional invariant sub-manifolds in the phase space of the general 2d Toda hierarchy. Indeed, functional reduction leaves a finite number of discrete indices in the ansatz for z as a function of w , while the string equation fixes the dependence of z on x , leaving finite number of degrees of freedom. These degrees of freedom are connected with the integration constants of system (12), which becomes finite-dimensional after functional reductions.

3.1 Polynomial reductions.

We begin with polynomial reductions of the 2d Toda hierarchy

$$z(w) = rw + \sum_{i=-N}^0 u_i w^i \quad (13)$$

$$\bar{z}(w^{-1}) = rw^{-1} + \sum_{i=0}^N \bar{u}_i w^i \quad (14)$$

constrained by the string equation (10).

The following proposition states the consistency of the polynomial reductions under the Toda flows:

Proposition 1 *If the string equation (10) holds, then (13), (14) is a finite dimensional subspace (of the phase space (5), (6)) invariant under the t_i and $\bar{t}_i$ 2d Toda flows, $0 < i < N + 2$ (7). This subspace has dimension $2N + 3$ with local coordinates chosen as the initial values of the solutions $r = r(x)$, $u_i = u_i(x)$, $\bar{u}_i = \bar{u}_i(x)$ to the reduced string equation (12).*

Proof. We must prove that z remains of the form (13), under the flows generated by $H_k, \bar{H}_k$ provided the string equation (10) holds. In other words, we have to show that the evolution does not change the highest and lowest degrees of the Laurent polynomial (13). The proof for $\bar{z}$ is analogous.

1. First we proceed with evolution of z generated by H_k . The lowest degree of w in z (13) is $-N$. Since

$$H_k = (z^k)_{>0} + 1/2(z^k)_0 = h_k(x)w^k + h_{k-1}(x)w^{k-1} + \dots + h_0(x) \quad (15)$$

is a Laurent polynomial of positive degree in w , the lowest degree of the bracket $\{z, H_k\}$ is not less than $-N$, as follows from (8). On the other hand, the complement $z^k - H_k$ of (15) is a polynomial in $1/w$ and so,

$$\{H_k, z\} = -\{z^k - H_k, z\}$$

is a Laurent polynomial whose highest degree in w does not exceed 1.

2. Unlike the H_k -flows, the form-invariance of z (13) under the flows generated by $\bar{H}_k$ requires extra restrictions on derivatives $\partial_x u_i, \partial_x \bar{u}_i$ (the string equation).

Since

$$\bar{H}_k = (\bar{z}^k)_- + 1/2(\bar{z}^k)_0$$

is a Laurent polynomial of non-positive degree, it conserves the highest degree of $z(w)$ under the evolution (7), but not the lowest degree, in general. Indeed,

$$\bar{H}_k = \bar{z}^k - ((\bar{z}^k)_+ + 1/2(\bar{z}^k)_0)$$

is a difference between $\bar{z}^k$ and a polynomial of nonnegative degree, so that

$$\{\bar{H}_k, z\} = \{\bar{z}^k, z\} - \{((\bar{z}^k)_+ + 1/2(\bar{z}^k)_0), z\}$$

The second bracket in the last expression preserves the lowest degree of z . Thus the lowest degree of $\{\bar{H}_k, z\}$ equals the lowest degree of $\{\bar{z}^k, z\} = k\bar{z}^{k-1}\{\bar{z}, z\}$. Imposing the extra restriction (10) (string equation), we see that the lowest degree does not exceed that of $\bar{z}^{k-1}$. Since the lowest degree of $\bar{z}$ is -1 and that of z is $-N$, it follows that $k < N + 2$.

Therefore, the form of (13) is preserved by $t_i, \bar{t}_i$ Toda flows with $i = 1, \dots, N + 1$, provided the string equation (10) holds. This completes the proof.

It is easy to prove the converse to Proposition 1, i.e.

Proposition 2 *The string equation (10) is a necessary condition for the existence of integrable polynomial flows of the form (9), (13), (14)*

Proof: This essentially uses the same arguments as those leading to the result of Proposition 1. For the first $N + 1$ z flows to be consistent with the polynomial reduction it is necessary that the Laurent expansion of $\{\bar{z}, z\}$ be of the form $\{\bar{z}, z\} = f(x) + \sum_{i>0} U_i(x)/w^i$. The corresponding condition for the $\bar{z}$ flows is $\{\bar{z}, z\} = \bar{f}(x) + \sum_{i>0} \bar{U}_i(x)w^i$. Both can be simultaneously satisfied only if

$$\{\bar{z}, z\} = f(x) = \bar{f}(x)$$

is independent of w . Differentiating with respect to t_i (or $\bar{t}_i$), i.e. along the flow lines, using the equations of motion (7), and the Jacobi identity we get

$$\frac{\partial f}{\partial t_i} = \{H_i, f\} = w \frac{\partial f}{\partial x} \frac{\partial H_i}{\partial w}.$$

Since $w \frac{\partial H_i}{\partial w}$ is w -dependent, while f is not, it follows that $\frac{\partial f}{\partial x} = \frac{\partial f}{\partial t_i} = 0$ and similarly $\frac{\partial f}{\partial \bar{t}_i} = 0$. Therefore f is a constant in x and also constant along the flows, implying (up to a constant scaling) that the string equation (10) holds.

In the following subsections we consider rational and logarithmic reductions as well as their Hamiltonian structures. Polynomial solutions can then be viewed as a special case. Nevertheless the corresponding limiting procedures are rather cumbersome and it is easier to consider the polynomial case separately. A derivation of the canonical linearizing variables in the polynomial case is given in Appendix B. This closely follows the analogous procedure given in Section 4 for the logarithmic case. Local canonical linearizing variables, which are constant and linear in the hierarchy times, turn out to be “Richardson’s harmonic moments” of the Hele-Shaw (Laplacian growth) problem when $z(w)$ and $\bar{z}(\bar{w})$ are identified with the conformal mapping from the w to the z plane. This is a simple proof of the relation (originally established in [12]) between the “times” of the 2dToda hierarchy and harmonic moments of the exterior Hele-Shaw problem.

An important consequence of the present section is the fact that the string equation (10) necessarily holds for reduced conformal mappings evolving under a sufficient number of 2dToda flows (i.e. sufficient for complete integrability). In the context of interface dynamics, *Darcy’s law thus turns out to be a corollary of integrability* for reduced systems.

3.2 Rational reductions.

We now consider a rational reduction for $z(w)$ and $\bar{z}(w)$, $N > 1$ given by

$$z(w) = \frac{q_{N+1}(w)}{p_N(w)} = \frac{rw^{N+1} + \sum_{i=0}^N a_i w^i}{w^N + \sum_{i=0}^{N-1} b_i w^i}, \quad (16)$$

$$\bar{z}(w^{-1}) = \frac{\bar{q}_{N+1}(w^{-1})}{\bar{p}_N(w^{-1})} = \frac{rw^{-(N+1)} + \sum_{i=0}^N \bar{a}_i w^{-i}}{w^{-N} + \sum_{i=0}^{N-1} \bar{b}_i w^{-i}}. \quad (17)$$

The following Lemma states the consistency such rational reductions under the t_i Toda flows for (16) and the $\bar{t}_i$ flows for (17).

Lemma 1 *The form of the function $z(w)$ in (16) is invariant under the t_i flows and the form (17) of $\bar{z}(w^{-1})$ is invariant under the $\bar{t}_i$ flows for any $i > 0$.*

Proof: Consider the flows generated by H_k . Since

$$H_k = (z^k)_+ + 1/2(z^k)_0 \quad (18)$$

is a polynomial of nonnegative degree in w , its complement $z^k - H_k$ is a polynomial in $1/w$. The Laurent expansion of brackets (8) $\{H_k, z\} = -\{z^k - H_k, z\}$ around infinity therefore has the following form

$$\{H_k, z\} = k_1(T)w + k_0(T) + k_{-1}(T)w^{-1} + \dots \quad (19)$$

However the Lax bracket (8) also implies that it is rational of the form

$$\{H_k, z\} = Q(w)/p_N(w)^2.$$

It follows from (19) that the degree of polynomial $Q(w)$ does not exceed $2N + 1$.

On the other hand,

$$\partial_{t_k} z = P(w)/p_N^2(w), \quad P(w) = p_N \partial_{t_k} q_{N+1} - q_{N+1} \partial_{t_k} p_N,$$

where the degree of the polynomial $P(w)$ also does not exceed $2N + 1$. Equating the coefficients of polynomial $P(w) - Q(w)$ to zero, we get a system of differential equations for r, a, b . The number of equations is $2N + 2$. Thus we get a compatible system of differential equations for $2N + 2$ unknowns r, a, b . A similar argument shows the form invariance of (17) under the $\bar{t}_k$ flows.

The consistency of the rational reduction defined in (17) under the t_i flows and (16) under $\bar{t}_i$ flows, respectively requires additional restrictions such as the string equation. But this only suffices to ensure form invariance under t_1 and $\bar{t}_1$ flows respectively.

Lemma 2 *The string equation (10) is a sufficient condition for $z(w)$ and $\bar{z}(w)$ ((16), (17)) to be form invariant under the first two-Toda flows. Flows generated by $H_k, \bar{H}_k$ with $k > 1$ are inconsistent with the rational reduction (16), (17).*

Remark: In the next section, we will show how to choose $4N + 2$ independent commuting flows preserving the rational form of z (16) and $\bar{z}$ (17), generated by vector fields that are infinite linear combinations of those generating the 2dToda hierarchy.

Proof: In order to prove that z remains of the same form (16) under the $\bar{t}_i$ Toda flows we would have to find a compatibility condition for a system of differential equations for $r(T), a(T), b(T)$, induced by (7).

Consider evolution of z along the flows generated by $\bar{H}_k$. Again we write

$$\partial z / \partial \bar{t}_k = S(w)/p_N^2(w), \quad S(w) = p_N \partial_{\bar{t}_k} q_{N+1} - q_{N+1} \partial_{\bar{t}_k} p_N$$

with $S(w)$ a polynomial in w of order at most $2N + 1$.

Since

$$\bar{H}_k = (\bar{z}^k)_- + 1/2(\bar{z}^k)_0 = \bar{h}_k w^{-k} + \bar{h}_{k-1} w^{-k+1} + \dots + \bar{h}_0 \quad (20)$$

is a Laurent polynomial of non-positive degree in w , the Laurent expansion of the corresponding bracket has the following form

$$\{\bar{H}_k, z\} = k_1(T)w + k_0(T) + k_{-1}(T)w^{-1} + \dots$$

The definition of the Lax bracket (8) and the expansion (20) implies that

$$\{\bar{H}_k, z\} = (U(w) + R(1/w))/p_N(w)^2$$

where $U(w)$ is a polynomial of degree at most $2N + 1$ and $R = w^{-1}(\eta_{k-1}(T)w^{-k+1} + \dots + \eta_0)$ is a polynomial in w^{-1} . If $R(1/w) = 0$, the number of equations will not exceed the number of unknowns. However, as we show below, the string equation does not imply the vanishing of R .

Introduce the new variable $y = w^{-1}$. The vanishing of R is equivalent to the following form of expansion in y

$$\{\bar{H}_k, z\} = f_0(T) + f_{-1}(T)y^{-1} + \dots \quad (21)$$

Consider this as a function of y and, as before, express $\bar{H}_k$ as a difference of $\bar{z}_k$ and a polynomial of nonnegative degree. Then

$$\begin{aligned} \{\bar{H}_k, z\} &= \{\bar{z}^k, z\} - \{((\bar{z}^k)_+ + 1/2(\bar{z}^k)_0), z\} = \\ &= k\bar{z}^{k-1}\{z, \bar{z}\} - \{((\bar{z}^k)_+ + 1/2(\bar{z}^k)_0), z\} \end{aligned} \quad (22)$$

Imposing the string equation (10) we see that the only flow that allows the above expansion (21) corresponds to $k = 1$. A similar argument shows that only the t_1 flow preserves the form (17) of $\bar{z}$ even if the string equation is imposed.

As seen from the above proof, there are only two flows generated by evolution operators of the form (9) compatible with the rational reduction. In fact, we should not expect more invariant flows associated

to the simple poles at $w = 0$ and $w = \infty$. In the polynomial case, the number of invariant flows was equal to the number of variables (polynomial coefficients), since one can associate n invariant flows to a pole of n th order, and the poles at zero and infinity are immovable.

However, below we introduce additional flows related to movable singularities of $z(w)$ and $\bar{z}(w^{-1})$ which do preserve the rational reduction (16), (17), extending results of Krichever for the KP case.

3.3 Additional flows for rational reductions of dKP hierarchy.

In this subsection we recall the theory of flows related to poles at a finite number of finite points applying an approach previously used by Krichever for dKP hierarchy. In the dispersionless limit, the dKP and 1dToda hierarchies are quite similar, while in 2dToda the existence of finite-dimensional reductions requires extra constraints.

Let us start by recalling [5], that on the phase space of extended Benney systems, i.e. rational dKP, there arise some new flows related to the pole structure of the corresponding maps. These additional flows were introduced by Krichever (see [9]).

Consider the partial fraction expansion under the above assumptions and the flows of the t_i type only. For the dKP hierarchy [5], [19] these reductions have the form

$$z(w) = w + u_0 + \sum_{\alpha=1}^N \frac{u_\alpha}{w - w_\alpha}. \quad (23)$$

The new flows associated to the poles are defined similarly to the polynomial case

$$\partial_{t_{k,\alpha}} z = \{B_{k,\alpha}, z\}, \quad \alpha = \infty, 1, 2, \dots \quad k = 0, 1, 2, \dots \quad (24)$$

The evolution functions associated with the pole structure of z are as follows :

$$B_{k,\infty} = (z(w)^k)_{\geq 0}, \quad ()_{\geq 0} := ()_+ + ()_0$$

for an immovable pole at infinity, while for each finite-distance pole there appear additional flows with evolution functions:

$$B_{k,\alpha} = (z(w)^k)_\alpha, \quad B_{0,\alpha} = \log(w_\alpha - w).$$

Here $z(w)_\alpha$ denotes the negative part of a formal Laurent expansion of $z(w)$ near the pole w_α :

$$f(w)_\alpha = \sum_{i>0} \frac{f_i}{(w - w_\alpha)^i}, \quad \text{if } f = \sum_{i \in \mathbb{Z}} \frac{f_i}{(w - w_\alpha)^i}. \quad (25)$$

These additional flows commute amongst themselves and with the ordinary 1dToda or dKP flows (associated with poles at infinity) .

3.4 Additional invariant flows of 2dToda system.

In the following, we will look at reductions of the 2dToda systems analogous to the above Benney-Krichever reductions of KP. We will thus choose the rational functions $z(w)$, $\bar{z}(1/w)$ appearing in (16) and (17) to have only simple poles and express these in the partial fraction form

$$\begin{aligned} z(w) &= rw + u_0 + \sum_{j=1}^n \frac{u_j}{w - w_j}, \\ \bar{z}(1/w) &= r/w + \bar{u}_0 + \sum_{j=1}^n \frac{\bar{u}_j}{1/w - \bar{w}_j}. \end{aligned} \quad (26)$$

Now, introduce a new set of $4n + 2$ evolution functions $H_{0,j}, H_{1,j}, \bar{H}_{0,j}, \bar{H}_{1,j}, \bar{H}_{1,\infty}$ defined by

$$H_{k,j}(w) = B_{k,j}(w) - \frac{1}{2}B_{k,j}(w=0), \quad \bar{H}_{k,j}(y) = \bar{B}_{k,j}(y) - \frac{1}{2}\bar{B}_{k,j}(y=0), \quad (27)$$

where $k = 0, 1$, $j = 1..n$ and

$$\begin{aligned} B_{1,\infty}(w) &= (z(w))_{\geq 0}, & B_{0,j} &= \log(r(w_j - w)), & B_{1,j}(w) &= (z(w))_j \\ \bar{B}_{1,\infty}(y) &= (\bar{z}(y))_{\geq 0}, & \bar{B}_{0,j} &= \log(r(\bar{w}_j - y)), & \bar{B}_{1,j}(y) &= (\bar{z}(y))_j, \quad y = 1/w. \end{aligned} \quad (28)$$

We denote the flow variables associated with $H_{0,j}$ as τ_{2j} , those associated with $H_{1,j}$ as τ_{2j-1} , and that associated with $H_{1,\infty}$ as τ_0 . The flow variables associated with $\bar{H}_{0,j}$, $\bar{H}_{1,j}$ and $\bar{H}_{0,\infty}$ are denoted $\bar{\tau}_{2j}$, $\bar{\tau}_{2j-1}$, $\bar{\tau}_0$ respectively. Correspondingly, we denote the evolution functions

$$\begin{aligned} h_0 &= H_{1,\infty} = rw + u_0/2, & \bar{h}_0 &= \bar{H}_{1,\infty} = r/w + \bar{u}_0/2, \\ h_{2j-1} &= H_{1,j} = \frac{u_j}{w-w_j} + \frac{u_j}{2w_j}, & \bar{h}_{2j-1} &= \bar{H}_{1,j} = \frac{\bar{u}_j}{\bar{w}_j-1/w} + \frac{1}{2}\bar{u}_j/\bar{w}_j, \\ h_{2j} &= H_{0,j} = \log(w_j - w) + 1/2 \log(r/w_j), & \bar{h}_{2j} &= \bar{H}_{0,j} = \log(\bar{w}_j - 1/w) + 1/2 \log(r/\bar{w}_j). \end{aligned} \quad (29)$$

Then the following proposition holds

Proposition 3 *The $4n + 2$ commuting Toda-Krichever flows*

$$\begin{aligned} \partial_{\tau_j} z &= \{h_j, z\}, & \partial_{\bar{\tau}_j} z &= \{\bar{h}_j, z\} \\ \partial_{\tau_j} \bar{z} &= \{h_j, \bar{z}\}, & \partial_{\bar{\tau}_j} \bar{z} &= \{\bar{h}_j, \bar{z}\}, \end{aligned} \quad j = 0..2n \quad (30)$$

preserve the rational form of $z(w)$ and $\bar{z}(1/w)$ (26) (or equally (17, 16)) provided the string equation (10) holds. The dimension of the reduced phase space equals $4n + 3$.

We defer the proof, since this follows as a limiting case of the more general logarithmic reduction introduced in the next subsection.

As in the polynomial case, the total number of form invariant flows preserving the string equation equals the dimension of the reduced phase space minus one. In what follows we show that these flows are Hamiltonian. Since the dimension of the phase space is odd and equals $4n + 3$, it is, in fact a Poisson manifold whose symplectic leaves have dimension $4n + 2$, which is exactly the number of commuting Toda-Krichever flows.

The above result holds for a more general setting. Below we introduce a logarithmic reduction of the 2dToda hierarchy and prove an analog of proposition 3 for logarithmic functions. Proposition 3 follows as a limiting case.

3.5 Logarithmic flows.

It is easier to prove the consistency of the rational reductions with the dynamics of the 2dToda system by first considering the more general logarithmic functions and then taking limits in which the branch points degenerate in pairs. Let us set

$$\begin{aligned} z &= r(x)w + u(x) + \sum_{i=1}^{n+1} a_i \log(w_i(x) - w), \\ \bar{z} &= r(x)w^{-1} + \bar{u}(x) + \sum_{i=1}^{n+1} \bar{a}_i \log(\bar{w}_i(x) - w^{-1}), \end{aligned} \quad (31)$$

where $a_i, \bar{a}_i$ are arbitrary constants, subject to the conditions

$$\sum_{i=1}^{n+1} a_i = 0, \quad \sum_{i=1}^{n+1} \bar{a}_i = 0, \quad (32)$$

which ensure absence of logarithmic singularities at infinity.

Introduce evolution functions as follows:

$$\begin{aligned} \mathcal{H}_0 &= r(x)w + \frac{1}{2}u(x), & \bar{\mathcal{H}}_0 &= \bar{r}(x)w^{-1} + \frac{1}{2}\bar{u}(x), \\ \mathcal{H}_j &= \log(w_j(x) - w) + \frac{1}{2} \log(r(x)/w_j(x)), & j &= 1..n+1. \\ \bar{\mathcal{H}}_j &= \log(\bar{w}_j(x) - w^{-1}) + \frac{1}{2} \log(r(x)/\bar{w}_j(x)), \end{aligned} \quad (33)$$

We then have

Proposition 4 *The string equation (10) is a necessary and sufficient condition for the existence of $2n+4$ commuting flows on the $2n+5$ dimensional space of functions of the form (31)*

$$\begin{aligned} \partial_{\tau_i} z &= \{\mathcal{H}_i, z\}, & \partial_{\bar{\tau}_i} z &= \{\bar{\mathcal{H}}_i, z\} \\ \partial_{\tau_i} \bar{z} &= \{\mathcal{H}_i, \bar{z}\}, & \partial_{\bar{\tau}_i} \bar{z} &= \{\bar{\mathcal{H}}_i, \bar{z}\} \end{aligned}, \quad i = 0..n+1..n+1 \quad (34)$$

In other words, the flows generated by (34), (33) are tangential to the manifold of such logarithmic functions if the string condition is imposed (and conversely). We therefore have $2n+4$ flows leaving invariant a $2n+5$ dimensional sub-manifold of the 2dToda system. These will be shown to be infinite linear combinations of vector fields generating the 2dToda flows (7). Equations (44, 45) will show how to express corresponding 2dToda times $t_i, \bar{t}_i$ in terms of the logarithmic flow parameters $\tau_i, \bar{\tau}_i$.

Proof

1. *Commutativity* : This will be shown to follow as Corollary 1 to Proposition 5 below.

2. *Consistency of (31) with equations of motion (34)*: As in the polynomial and rational cases the consistency of the $\partial_{\tau_i} z$ and $\partial_{\bar{\tau}_i} \bar{z}$ equations with the logarithmic reduction (31) follows from the fact that the structure of the Lax-Poisson brackets $\{\mathcal{H}_i, z\}, \{\bar{\mathcal{H}}_i, \bar{z}\}$ is identical to the infinitesimal deformations in z and $\bar{z}$ induced by ones in the functions $r, \bar{u}, u, \bar{w}_i, w_i$. However the consistency of the $\partial_{\tau_i} \bar{z}$ and $\partial_{\bar{\tau}_i} z$ equations with the logarithmic reduction requires an extra constraint, the string equation. First, we prove this for $\partial_{\bar{\tau}_i} z$.

Differentiating z in (31) with respect to $\bar{\tau}_i$ using the equation of motion (34), we get

$$\partial_{\bar{\tau}_i} z = w \partial_{\bar{\tau}_i} r + \partial_{\bar{\tau}_i} u + \sum_{j=1}^{n+1} \frac{a_i \partial_{\bar{\tau}_i} w_j}{w - w_j} = \{\bar{\mathcal{H}}_i, z\}. \quad (35)$$

The left-hand side of (35) may contain any terms that are linear in w at $w = \infty$ and simple poles at $w = w_j, j = 1..n+1$. Since the Lax-Poisson bracket (8) is a bi-derivation, the singularities in w that may occur in $\{\bar{\mathcal{H}}_i, z\}$ consist either of simple poles at $w = w_j, j = 1..n+1$, linear terms at $w = \infty$ or a simple pole at $w = 1/\bar{w}_i$. We show that the latter is absent.

Note that $\bar{z}$ can be represented as a sum over $\bar{\mathcal{H}}_i, i = 0..n+1$ plus a w -independent function $f(x)$;

$$\bar{z} = \sum_{i=0}^{n+1} \bar{a}_i \bar{\mathcal{H}}_i + f(x), \quad (36)$$

where $\bar{a}_0 := 1$, $f(x) = \frac{1}{2} \left(\bar{u}(x) - \sum_{i=1}^{n+1} a_i \log \bar{w}_i(x) \right)$. Now, using (36) we get

$$\bar{a}_i \{\bar{\mathcal{H}}_i, z\} = \{\bar{z} - \sum_{j \neq i} \bar{a}_j \bar{\mathcal{H}}_j - f(x), z\} = \{\bar{z}, z\} - \sum_{j \neq i} \bar{a}_j \{\bar{\mathcal{H}}_j, z\} - \{f(x), z\}. \quad (37)$$

Since $f(x)$ is independent of w , the last term in (37) contains only linear terms in w plus simple poles at $w_j, j = 1..n+1$. The term $\sum_{j \neq i} \bar{a}_j \{\bar{\mathcal{H}}_j, z\}$ could, in principle, contain poles at $w = 1/\bar{w}_j$ for $j \neq i$, but these must cancel since $\{\bar{\mathcal{H}}_i, z\}$ contains no such poles. Furthermore, this term contains no pole at $w = 1/\bar{w}_i$, because $\bar{\mathcal{H}}_i$ is omitted in the sum. Since the string equation (10) holds by assumption, we see that the rhs of (35) only contains terms of the type induced by infinitesimal deformations in the functions r, u, w_j , showing that the $\partial_{\bar{\tau}_i} z$ equation, together with the string equation, is compatible with the reduction (31). The proof for the $\partial_{\tau_i} \bar{z}$ equation is similar.

The above proves the sufficiency of the string equation for the validity of Proposition 4. The necessity is proved similarly to Proposition 2. By equation (37), if the rational reduction (31) is preserved under the $\bar{\tau}_i$ flows, $\{\bar{z}, z\}$ can have no pole at $w = 1/\bar{w}_i$ for any i or at zero. Similarly to preserve (31) under the τ_i flows it can have no poles at $w_i, i = 1..n+1$ or ∞ . But since the only possible poles in $\{\bar{z}, z\}$ are at these points, we conclude that $\{\bar{z}, z\}$ is constant in w and hence can only be a function (say q) of $x, \tau, \bar{\tau}$

$$\{\bar{z}, z\} = q$$

Differentiating the last equation with respect to τ_i (or $\bar{\tau}_i$) using the equation of motion (34), the Jacobi identity and the definition of the Lax-Poisson bracket we get

$$\frac{\partial q}{\partial \tau_i} = w \frac{\partial q}{\partial x} \frac{\partial \mathcal{H}_i}{\partial w}.$$

Since $w \frac{\partial \mathcal{H}_i}{\partial w}$ is w -dependent, while q is not, it follows that $\frac{\partial q}{\partial x} = \frac{\partial q}{\partial \tau_i} = 0$ (similarly $\frac{\partial q}{\partial \bar{\tau}_i} = 0$). Therefore $q = \text{const}$ and the string equation (10) holds.

The demonstration that the reduced phase space is of dimension $2n + 5$ is similar to the polynomial case. We view the string equation (10) together with the logarithmic reduction (31) as a set of $(2n + 5)$ first order ODE's for the $(2n + 5)$ functions $r(x), u(x), \bar{u}(x), w_i(x), \bar{w}_i(x), i = 1..n + 1$

$$\begin{aligned} \frac{dr}{dx} &= R(r, u, \bar{u}, w_1, \dots, \bar{w}_1, \dots), & \frac{du}{dx} &= U(r, u, \bar{u}, w_1, \dots, \bar{w}_1, \dots), & \frac{d\bar{u}}{dx} &= \bar{U}(r, u, \bar{u}, w_1, \dots, \bar{w}_1, \dots) \\ \frac{dw_i}{dx} &= W_i(r, u, \bar{u}, w_1, \dots, \bar{w}_1, \dots), & \frac{d\bar{w}_i}{dx} &= \bar{W}_i(r, u, \bar{u}, w_1, \dots, \bar{w}_1, \dots), & i &= 1..n + 1 \end{aligned}$$

which, at least locally, determine the dependence of these functions uniquely in terms of their values $r_0, u_0, \bar{u}_0, w_{i0}, \bar{w}_{i0}, i = 1..n + 1$ at some initial value $x = x_0$

$$r = r(r_0, u_0, \bar{u}_0, w_{1,0}, \dots, \bar{w}_{1,0}, \dots, x), \quad u = u(r_0, u_0, \bar{u}_0, w_{1,0}, \dots, \bar{w}_{1,0}, \dots, x), \quad \bar{u} = \bar{u}(r_0, u_0, \bar{u}_0, w_{1,0}, \dots, \bar{w}_{1,0}, \dots, x)$$

$$w_i = w_i(r_0, u_0, \bar{u}_0, w_{1,0}, \dots, \bar{w}_{1,0}, \dots, x), \quad \bar{w}_i = \bar{w}_i(r_0, u_0, \bar{u}_0, w_{1,0}, \dots, \bar{w}_{1,0}, \dots, x), \quad i = 1..n + 1$$

The $2n + 5$ initial values $r_0, u_0, \bar{u}_0, w_{i0}, \bar{w}_{i0}, i = 1..n + 1$ may be viewed as coordinates of the reduced phase space.

Proof of Proposition 3: This follows from taking limits in which the logarithmic branch points w_{2i}, w_{2i-1} coalesce in pairs.

Setting

$$\begin{aligned} a_{2i-1} &= 1/\epsilon, & a_{2i} &= -1/\epsilon & w_{2i} &= w_{2i-1} + \epsilon u_i \\ \bar{a}_{2i-1} &= 1/\epsilon, & \bar{a}_{2i} &= -1/\epsilon & \bar{w}_{2i} &= \bar{w}_{2i-1} + \epsilon \bar{u}_i \end{aligned} \tag{38}$$

in (31) we get (26) in the $\epsilon \rightarrow 0$ limit.

The evolution functions (29), generating flows on the space of rational reduction (26), are then obtained as follows:

$$\begin{aligned} h_0 &= \lim_{\epsilon=0} \mathcal{H}_0, & \bar{h}_0 &= \lim_{\epsilon=0} \bar{\mathcal{H}}_0 \\ h_{2i-1} &= \lim_{\epsilon=0} \frac{1}{\epsilon} (\mathcal{H}_{2i} - \mathcal{H}_{2i-1}), & \bar{h}_{2i} &= \lim_{\epsilon=0} \frac{1}{\epsilon} (\bar{\mathcal{H}}_{2i} - \bar{\mathcal{H}}_{2i-1}) \\ h_{2i} &= \lim_{\epsilon=0} \mathcal{H}_{2i-1}, & \bar{h}_{2i-1} &= \lim_{\epsilon=0} \bar{\mathcal{H}}_{2i-1}. \end{aligned}$$

We thus obtain evolution functions for the rational reductions, where $z, \bar{z}$ have simple poles only, as a limiting degenerating logarithmic case. One may deduce structures related to general rational reductions by degenerating arbitrary numbers of logarithmic singularities at different finite points as well as at infinity.

4 Poisson structure of logarithmic reductions of the 2Toda hierarchy.

In this section we study the Poisson structure of logarithmic reductions of the 2dToda hierarchy, which will be shown to define finite-dimensional completely integrable systems. We find explicit expressions leading to a canonical Hamiltonian structure on the phase space of rational reductions of the 2dToda system.

The Poisson structure of rational reductions of the 1dToda hierarchy (which are infinite-dimensional, having no string equation constraint) is described in Appendix C. It is related to the Poisson structure of the Benney system considered in [14]

4.1 Hamiltonians, Action-Angle variables.

Let us now introduce the following $2n+5$ functions on the reduced phase space defined by (31) with $z, \bar{z}$ solutions to (10), extended by the auxiliary variable x

$$r, u, \bar{u}, w_1, \dots, w_{n+1}, \bar{w}_1, \dots, \bar{w}_{n+1} \rightarrow Q, I_0, I_1, \dots, I_{n+1}, \bar{I}_0, \bar{I}_1, \dots, \bar{I}_{n+1}, \quad (39)$$

where the new variables are defined as follows

$$\begin{aligned} I_0 &= \frac{1}{2i\pi} \oint_{\infty} \bar{z} d \ln z, & \bar{I}_0 &= \frac{1}{2i\pi} \oint_0 z d \ln \bar{z}, \\ I_j &= \frac{1}{2i\pi} \oint_{w_j} \bar{z} dz, & \bar{I}_j &= \frac{1}{2i\pi} \oint_{1/\bar{w}_j} z d\bar{z}, \quad j = 1..n+1, \\ Q &= \frac{1}{4i\pi} \sum_{i=0}^{n+1} \oint_{1/\bar{w}_i} z d\bar{z} + \oint_{w_i} \bar{z} dz. \end{aligned}$$

Using (31), we may evaluate the contour integrals in the last equation, expressing these explicitly in terms of the old parametrization $r, u, \bar{u}, w_j, \bar{w}_j, j = 1..n+1$

$$\begin{aligned} I_0 &= \bar{z}(1/w=0) = \bar{u} + \sum_{j=1}^{n+1} \bar{a}_j \ln(\bar{w}_j), & \bar{I}_0 &= z(0) = u + \sum_{j=1}^{n+1} a_j \ln(w_j), \\ I_j &= a_j \bar{z}(w_j^{-1}) = a_j \left(r w_j^{-1} + \bar{u} + \sum_{k=1}^{n+1} \bar{a}_k \ln(\bar{w}_k - w_j^{-1}) \right), \\ \bar{I}_j &= \bar{a}_j z(\bar{w}_j^{-1}) = \bar{a}_j \left(r \bar{w}_j^{-1} + u + \sum_{k=1}^{n+1} a_k \ln(w_k - \bar{w}_j^{-1}) \right), \quad j = 1..n+1, \end{aligned} \quad (40)$$

$$\begin{aligned} Q &= \frac{1}{2} r \left(\left(\frac{\partial z(w)}{\partial w} \right)_{w=0} + \left(\frac{\partial \bar{z}(1/w)}{\partial (1/w)} \right)_{1/w=0} \right) - \frac{1}{2} \sum_{j=1}^{n+1} (I_j + \bar{I}_j) \\ &= r^2 - \frac{1}{2} \sum_{j=1}^{n+1} \left(r \left(\frac{a_j}{w_j} + \frac{\bar{a}_j}{\bar{w}_j} \right) + I_j + \bar{I}_j \right). \end{aligned} \quad (41)$$

Proposition 5 *The functions $I_k, \bar{I}_k, Q$ (40, 41) are linearizing variables of the system (34) satisfying*

$$\begin{aligned} \partial_{\tau_j} Q &= 0, & \partial_{\bar{\tau}_j} Q &= 0, \\ \partial_{\tau_j} I_k &= \delta_{jk}, & \partial_{\bar{\tau}_j} I_k &= 0, \\ \partial_{\tau_j} \bar{I}_k &= 0, & \partial_{\bar{\tau}_j} \bar{I}_k &= -\delta_{jk}. \end{aligned} \quad (42)$$

Proof: We first calculate derivatives of each I_k with respect to the times $\tau_j, j = 0..n+1$. Applying integration by parts, we get

$$2\pi i \frac{\partial I_k}{\partial \tau_j} = \frac{\partial}{\partial \tau_j} \oint_{w_k} \bar{z} \frac{\partial z}{\partial w} dw = \oint_{w_k} \frac{\partial}{\partial w} \left(\bar{z} \frac{\partial z}{\partial \tau_j} \right) dw + \oint_{w_k} \left(\frac{\partial \bar{z}}{\partial \tau_j} \frac{\partial z}{\partial w} - \frac{\partial z}{\partial \tau_j} \frac{\partial \bar{z}}{\partial w} \right) dw.$$

By the equations of motion (34) and the definition of the Lax-Poisson brackets (8) this equals

$$\begin{aligned} &\oint_{w_k} \frac{\partial}{\partial w} \left(\bar{z} \frac{\partial z}{\partial \tau_j} \right) dw + \oint_{w_k} \left(\{ \mathcal{H}_j, \bar{z} \} \frac{\partial z}{\partial w} - \{ \mathcal{H}_j, z \} \frac{\partial \bar{z}}{\partial w} \right) dw \\ &= \oint_{w_k} \frac{\partial}{\partial w} \left(\bar{z} \frac{\partial z}{\partial \tau_j} \right) dw + \oint_{w_k} \frac{\partial \mathcal{H}_j}{\partial w} w \left(\frac{\partial z}{\partial w} \frac{\partial \bar{z}}{\partial x} - \frac{\partial z}{\partial x} \frac{\partial \bar{z}}{\partial w} \right) dw, \end{aligned}$$

which, by the string equation (10) reduces to:

$$\oint_{w_k} \frac{\partial}{\partial w} \left(\bar{z} \frac{\partial z}{\partial \tau_j} \right) dw + \oint_{w_k} \frac{\partial \mathcal{H}_j}{\partial w} dw = 2\pi i \delta_{kj}, \quad \delta_{jk} = \begin{cases} 1, & j = k \\ 0, & j \neq k. \end{cases}$$

The first term in the lhs vanishes, because $\bar{z} \frac{\partial z}{\partial \tau_j}$ is univalent in a neighbourhood of $w_k, k = 1..n+1$ (and at ∞ due to (32)) and the remaining integral is evaluated by substituting expression (33) for $\mathcal{H}_j$. We have thus proved that

$$\frac{\partial I_i}{\partial \tau_j} = \delta_{ij}.$$

The rest of the proposition is proved by similar computations.

Corollary 1 *The vector fields defining (34) commute.*

Proof: Since $z, \bar{z}$ are completely determined by the new coordinates $Q, I, \bar{I}$, infinitesimal deformations of the former under the τ_i flows can be expressed via the chain rule through the infinitesimal deformations of the latter. It follows from Proposition 5 that

$$\frac{\delta}{\delta \tau_i} \begin{pmatrix} z \\ \bar{z} \end{pmatrix} = \frac{\partial}{\partial I_i} \begin{pmatrix} z \\ \bar{z} \end{pmatrix}, \quad \frac{\delta}{\delta \bar{\tau}_i} \begin{pmatrix} z \\ \bar{z} \end{pmatrix} = \frac{\partial}{\partial \bar{I}_i} \begin{pmatrix} z \\ \bar{z} \end{pmatrix},$$

so we have

$$\left(\frac{\delta}{\delta \tau_i} \frac{\delta}{\delta \tau_j} - \frac{\delta}{\delta \tau_j} \frac{\delta}{\delta \tau_i} \right) \begin{pmatrix} z \\ \bar{z} \end{pmatrix} = \left(\frac{\partial}{\partial I_i} \frac{\partial}{\partial I_j} - \frac{\partial}{\partial I_j} \frac{\partial}{\partial I_i} \right) \begin{pmatrix} z \\ \bar{z} \end{pmatrix} = 0.$$

The commutativity of the other flows are seen to similarly follow from Proposition 5.

Choosing the Poisson structure for equations (42) in a canonical way

$$\{I_i, \bar{I}_j\}_p = \delta_{ij}, \quad \{I_i, I_j\}_p = \{\bar{I}_i, \bar{I}_j\}_p = 0, \quad \{Q, \bar{I}_j\}_p = \{Q, I_j\}_p = 0, \quad (43)$$

we may interpret $I, \bar{I}$ as canonical linearizing variables and Q as a Casimir invariant. The equations of motion induced by the evolution operators $\mathcal{H}_i$ or $\bar{\mathcal{H}}_i$ are seen to be Hamiltonian and generated by the Hamiltonians $\bar{I}_i, I_i$.

The Poisson brackets $\{, \}_p$ in (43) are different from $\{, \}$ in (8); they define a Poisson structure on the finite dimensional space of logarithmic reductions of the phase space satisfying the string equation, while the latter (the Lax-Poisson bracket) is a dispersionless limit of the commutator.

Remark: Linearizing coordinates similar to (40) appeared in [11] in connection with the Laplacian Growth problem as integrals of the Laplacian Growth (string) equation, without the introduction of the compatible τ_i and $\bar{\tau}_i$ flows considered here.

The following proposition makes explicit the fact that the string equation implies the finite dimensionality of the reduced phase space.

Proposition 6 *The functions (40) form a set of integrals of the string equation (10). The string equation (10) implies*

$$\frac{\partial I_i}{\partial x} = \frac{\partial \bar{I}_i}{\partial x} = 0, \quad \frac{\partial Q}{\partial x} = 1.$$

Equivalently, this may be integrated to

$$Q - x = c_0, \quad I_i = c_i, \quad \bar{I}_i = \bar{c}_i, \quad i = 0..n+1,$$

where $c_i, \bar{c}_i$ are constants in x , which may be interpreted as coordinates on the $2n+5$ dimensional reduced phase space.

Proof: This is similar to the proof of proposition (5); one simply differentiates with respect to x and evaluates corresponding residues.

As mentioned in the Introduction, the Laplacian growth problem is recovered by identifying $\bar{z}$ as the complex conjugate of z . As seen from Proposition 6, the Casimir Q , which is proportional to the area of $\mathcal{D}_+$ grows with unit speed in physical time x , while $I_k, \bar{I}_k$ are functions of the harmonic moments of the boundary curve.

To obtain similar results for the rational case we take a limit as in (38).

Corollary 2 *Let $z, \bar{z}$ be of the rational reduction form (26). Then the following $2n+3$ quantities*

$$\begin{aligned} I_0 &= \bar{u}_0 - \sum_{i=1}^n \bar{u}_i / \bar{w}_i, & \bar{I}_0 &= u_0 - \sum_{i=1}^n u_i / w_i \\ I_{2i-1} &= r w_i^{-1} + \bar{u}_0 + \sum_{j=1}^n \frac{\bar{u}_j}{1/\bar{w}_i - \bar{w}_j}, & \bar{I}_{2i-1} &= r \bar{w}_i^{-1} + u_0 + \sum_{j=1}^n \frac{u_j}{1/\bar{w}_i - w_j}, \quad i = 1..n \\ I_{2i} &= \left(r - \sum_{j=1}^n \frac{\bar{u}_j}{(1/\bar{w}_i - \bar{w}_j)^2} \right) \frac{u_i}{w_i^2}, & \bar{I}_{2i} &= \left(r - \sum_{j=1}^n \frac{u_j}{(1/\bar{w}_i - w_j)^2} \right) \frac{\bar{u}_i}{\bar{w}_i^2} \end{aligned}$$

$$Q = r^2 - \frac{1}{2} \sum_{i=1}^n \left(r \left(\frac{\bar{u}_i}{\bar{w}_i^2} + \frac{u_i}{w_i^2} \right) + \bar{I}_{2i} + I_{2i} \right)$$

the linearizing canonical variables for the rational reduction, i.e. variables in terms of which the equations of motion (30) for the rational reduction (26) have the canonical form (42).

The logarithmic or rational flows are infinite linear combinations of the 2dToda flows (7). As shown in Appendix B, harmonic moments of the conformal mapping $z(w)$ are linear in the 2dToda times. Therefore to express $t_i, \bar{t}_i, i = 1, 2..$ through $\tau_i, \bar{\tau}_i, i = 0..n + 1$ one has to evaluate the integrals

$$M_k(z, \bar{z}) = \frac{1}{2k\pi i} \oint_{\infty} \bar{z} z^{-k} \frac{\partial z}{\partial w} dw, \quad \bar{M}_k(z, \bar{z}) = \frac{1}{2k\pi i} \oint_0 z \bar{z}^{-k} \frac{\partial \bar{z}}{\partial w} dw, \quad (44)$$

which are functions of coordinates $z, \bar{z}$ of the reduced phase space (31), and then express $z, \bar{z}$ through $Q, I_i, \bar{I}_i, i = 0..n + 1$. Then

$$M_k(Q, I, \bar{I}) = M_k(z(Q, I, \bar{I}), \bar{z}(Q, I, \bar{I})), \quad \bar{M}_k(Q, I, \bar{I}) = \bar{M}_k(z(Q, I, \bar{I}), \bar{z}(Q, I, \bar{I})) \quad (45)$$

Since, modulo integration constants

$$M_i = t_i, \quad \bar{M}_i = \bar{t}_i, \quad i > 0,$$

$$I_i = \tau_i, \quad \bar{I}_i = \bar{\tau}_i, \quad i = 0..n + 1$$

and $Q = x$, it follows that equation (45) expresses $t_i, \bar{t}_i$ in terms of $\tau_i, \bar{\tau}_i$.

5 Conclusions.

We have derived consistent finite dimensional logarithmic and rational reductions of certain flows related to the 2dToda hierarchy. The requirement for the consistency of these reductions was the string equation. Since the latter may also be viewed as the Galin-Polubarinova equation for the Hele-Shaw (Laplacian growth) process, this also established the latter as a compatible constraint for such reduced 2dToda flows.

More generally, it would be of interest to determine all finite dimensional reductions of the 2dToda hierarchy that imply fulfilment of the Darcy law in various forms.

Also, it would be useful to study integrable systems connected to various kinds of Hele-Shaw flows. This could include boundary conditions which are more general than a point sink at infinity. For instance, one can consider an exterior problem with a steady and uniform viscous flow at infinity. This can describe the evolution of a bubble surrounded by a viscous liquid moving through a wide channel, or a bubble within a viscous flow generated by a source and a sink of equal magnitude separated by a large distance. In such models, the constraint on $z, \bar{z}$ (5, 6) has the form

$$\{z(w, x), \bar{z}(w^{-1}, x)\} = (1/w + w)r(x)$$

This is an exterior analog of the interior problem with a dipole source inside the domain. In contrast to a monopole sink, this implies that the first harmonic moment changes with the time x , while the others (including area) remain unchanged. In general any combination of n -pole sources could be considered:

$$\{z(w, x), \bar{z}(w^{-1}, x)\} = f(x, w)$$

where f is defined by the asymptotics of the hydrodynamic potential.

6 Appendix A: String equation and Darcy law

We recall here the derivation of the Galin-Polubarinova equation. Consider evolution of the exterior $\mathcal{D}_-$ of a simply-connected planar domain bounded by an analytic curve on z -plane. The curve is an image of

the unit circle in the w plane under a conformal mapping (3). As stated in the introduction the pressure field $P(X, Y)$ is constant in the interior domain $\mathcal{D}_+$ and at the boundary $\partial\mathcal{D}_+$, so that at $\partial\mathcal{D}_+$

$$0 = dP/dx = \partial P/\partial x + v_n \nabla P = \partial P/\partial x - (\nabla P)^2,$$

where we used Darcy law (2) and the pressure gradient is taken in the exterior vicinity of the curve. Since P is harmonic in $\mathcal{D}_-$ (cf (1)), it is the real part of a homomorphic function (hydrodynamic potential) $P = \text{Re}\phi(z)$. It follows that

$$0 = \text{Re} \left(\frac{\partial \phi}{\partial x} - \frac{\partial \phi}{\partial z} \overline{\left(\frac{\partial \phi}{\partial z} \right)} \right).$$

The hydrodynamic potential can be chosen as $\phi(z, x) = 1/2\pi \log w(z, x)$ due to the logarithmic asymptotic in (1). Then

$$0 = \text{Re} \left(\frac{1}{w(z, x)} \frac{\partial w(z, x)}{\partial x} - \frac{\partial w(z, x)}{\partial z} \overline{\left(\frac{\partial w(z, x)}{\partial z} \right)} \right)$$

where $w(z, x)$ stands for the map inverse to $z(w, x)$. Using the facts that $z = z(w(z, x), x)$; i.e.,

$$\frac{\partial z(w, x)}{\partial w} \frac{\partial w(z, x)}{\partial x} + \frac{\partial z(w, x)}{\partial x} = 0, \quad \frac{\partial w(z, x)}{\partial z} = 1 / \frac{\partial z(w, x)}{\partial w},$$

and $\bar{w} = 1/w$ on $\partial\mathcal{D}_+$, we arrive at (4).

7 Appendix B: Poisson structure of polynomial reductions

Following steps similar to the proof of Proposition 5, one arrives at the following result for polynomial reductions (13), (14), (9) of the 2dToda system.

Proposition 7 *The following quantities*

$$M_k = \frac{1}{2i\pi k} \oint_{\infty} \bar{z} z^{-k} dz, \quad \bar{M}_i = \frac{1}{2i\pi k} \oint_0 z \bar{z}^{-k} d\bar{z},$$

$$Q = \frac{1}{4i\pi} \left(\oint_{\infty} \bar{z} dz - \oint_0 z d\bar{z} \right)$$

are canonical linearizing variables of the (polynomially) reduced 2dToda system (13), (14), (9), (7):

$$M_k(z, \bar{z}) = t_k + \text{const}, \quad \bar{M}_k(z, \bar{z}) = \bar{t}_k + \text{const}, \quad Q(z, \bar{z}) = x + \text{const}.$$

Solutions to the string equation (10) are then defined by the set of algebraic equations

$$Q(z, \bar{z}) = x + c_0, \quad M_k(z, \bar{z}) = c_k, \quad \bar{M}_k(z, \bar{z}) = \bar{c}_k,$$

where $c_k, \bar{c}_k$ are arbitrary constants.

Proof: Evaluating $\frac{\partial M_i}{\partial t_k}, \frac{\partial M_j}{\partial t_k}$ by arguments similarly to those of proposition 5, we get

$$\frac{\partial M_j}{\partial t_k} = -\frac{1}{2i\pi j} \oint_{\infty} \frac{\partial H_k}{\partial w} z^{-i} dw, \quad \frac{\partial M_i}{\partial t_j} = \frac{1}{2i\pi j} \oint_{\infty} \frac{\partial \bar{H}_k}{\partial w} z^{-i} dw.$$

Using the fact that

$$H_k = (z(w)^k)_+ + \frac{1}{2}(z(w)^k)_0 = z(w)^k - (z(w)^k)_- - \frac{1}{2}(z(w)^k)_0$$

and the negative part of the Laurent expansion does not contribute to the integral, we obtain

$$\frac{\partial M_j}{\partial t_k} = \frac{1}{2i\pi j} \oint_{\infty} z(w)^{-i} \frac{\partial z(w)^j}{\partial w} dw = \delta_{kj}.$$

Using the fact that the evolution functions $\bar{H}_j = (\bar{z}^j)_- + 1/2(\bar{z}^j)_0$ generating the $\bar{\tau}_i$ flows, are polynomials of nonnegative degrees in $1/w$, we obtain

$$\frac{\partial M_i}{\partial \bar{t}_j} = 0.$$

In the context of the Hele-Shaw problem, the first $N + 1$ Richardson moments [16] equal the first $N + 1$ 2dToda times. Indeed, in the Hele-Shaw problem $z(w)$ has the meaning of a conformal mapping from w to the z plane. In the exterior problem, the exterior of the unit circle (i.e. domain $|w| > 1$) is mapped to the exterior $\mathcal{D}_-$ of the boundary curve $z(w)$. The exterior harmonic moments are, by Green's theorem

$$\frac{1}{2j\pi i} \int_{|w|>1} z^{-j} dz d\bar{z} = \frac{1}{2j\pi i} \oint_{|w|=1} \bar{z}(1/w) z(w)^{-j} dz.$$

Since the mapping $z(w)$ is univalent in $\mathcal{D}_-$, all zeros of $z(w)$ are located in $\mathcal{D}_+$ and we can move the integration contour to infinity if $i \geq 1$. Thus, we get

$$\frac{1}{2j\pi i} \oint_{\infty} \bar{z}(1/w) z(w)^{-j} dz = M_i$$

which, by proposition 7 equals t_i modulo an integration constant.

8 Appendix C: Poisson structure of 1dToda hierarchy.

In this section we consider the Poisson structure of rational reductions of 1dToda system.

Recall that for the 1dToda system one takes into account only the t_i flows.

$$\partial_{t_i} z = \{H_i, z\}, \quad H_i = (z(w)^i)_+ + 1/2(z(w)^i)_0, \quad i = 0.. \infty \quad (46)$$

This system is bi-Hamiltonian (for general information e.g. see [3]) with two (linear and quadratic) compatible Poisson structures. For generic Toda system (5), the dispersionless linear Poisson brackets for the "field variables" $u_i, i = 1.. \infty$ (5) have the following form [2]

$$\{u_n(x), u_m(y)\}_1 = 2(c_n + c_m - 1)[(n + m)u_{n+m}(x)\delta'(x - y) + mu'_{n+m}(x)\delta(x - y)] \quad (47)$$

where

$$c_k = \begin{cases} 1 & , \text{ if } k > 0 \\ 1/2 & , \text{ if } k = 0 \\ 0 & , \text{ if } k < 0 \end{cases}$$

while the quadratic brackets are

$$\begin{aligned} \{u_n(x), u_m(y)\}_2 = & \left[\frac{1}{2}(n - m)u_n(x)u'_m(x) + \left(\sum_{k=1}^{1-n} (n - m + k)u_{n+k}(x)u'_{m-k}(x) + \right. \right. \\ & \left. \left. + ku'_{n+k}(x)u_{m-k}(x) \right) \right] \delta(x - y) + \left[1/2(n - m)u_n u_m + \right. \\ & \left. + \left(\sum_{k=1}^{1-n} (n - m + 2k)u_{n+k}u_{m-k} \right) \right] \delta(x - y). \end{aligned} \quad (48)$$

As seen from the lemma 1, the rational functions

$$z(w) = \frac{q_{N+1}(w)}{p_N(w)} = \frac{w^{N+1} + \sum_{i=0}^N a_i w^i}{\sum_{i=0}^N b_i w^i} \quad (49)$$

are form-invariant under all the 1dToda flows (46), without any extra restriction (e.g. no string equation is needed).

We obtain corresponding Poisson structures for the variables a_i, b_i , by using result (48), expressing u_i in terms of $a_i, b_i, i = 0..N$. Both the linear (47) and the quadratic (48) Poisson structures lead to quadratic brackets for a_i, b_i . Namely, the second Poisson structure for (49) becomes

$$\{a_k(x), a_l(y)\}_2 = \left[\sum_{n=1} (l+n-k) a_{k-n}(x) a_{l+n}(y) + n a_{k-n}(y) a_{l+n}(x) \right. \\ \left. + (l-N-1) a_k(x) a_l(y) \right] \delta'(x-y), \quad (50)$$

$$\{b_k(x), b_l(y)\}_2 = \left[\sum_{n=1} (k-l-n) b_{k-n}(x) b_{l+n}(y) - n b_{k-n}(y) b_{l+n}(x) \right. \\ \left. + \frac{k-l}{2} b_k(x) b_l(y) \right] \delta'(x-y), \quad (51)$$

$$\{a_k(x), b_l(y)\}_2 = \frac{k-N-1}{2} a_k(x) b_l(y) \delta'(x-y). \quad (52)$$

The first Poisson structure can be obtained from (50 - 52) with the help of a shift by a constant

$$a_i \rightarrow a_i + \lambda b_i, \quad z(w, x) \rightarrow z(w, x) + \lambda$$

and using a bi-Hamiltonian nature of (47), (48).

$$\{a_k(x), a_l(y)\}_1 = \left[\sum_{n=1} (k-l-n) (a_{k-n}(x) b_{l+n}(y) + b_{k-n}(x) a_{l+n}(y)) - \right. \\ \left. - n (a_{k-n}(y) b_{l+n}(x) + b_{k-n}(y) a_{l+n}(x)) \right] + \frac{N+1-l}{2} b_k(x) a_l(y) + \\ + \frac{k+N+1-2l}{2} a_k(x) b_l(y) \delta'(x-y), \quad (53)$$

$$\{a_k(x), b_l(y)\}_1 = \left[\sum_{n=1} ((k-l-n) b_{k-n}(x) b_{l+n}(y) - n b_{k-n}(y) b_{l+n}(x)) + \right. \\ \left. + \frac{N+1-l}{2} b_k(x) b_l(y) \right] \delta'(x-y), \quad (54)$$

$$\{b_k(x), b_l(y)\}_1 = 0. \quad (55)$$

In all the above expressions $a_{N+1} = 1$ and $a_i = 0$ if i goes beyond the range $i = 0..N+1$ (and $b_j = 0$ if $j \neq 0..N$).

These brackets form a bi-Hamiltonian structure for rational reductions of 1dToda hierarchy:

$$\partial_{t_i} z = \{H_i, z\}_1 = \{H_{i-1}, z\}_2 \quad (56)$$

with the following Hamiltonians:

$$H_i = \frac{1}{i+1} \int (z^{i+1}(x))_0 dx \quad (57)$$

Acknowledgements

The authors would like to acknowledge helpful discussions and usefull information received from B.Dubrovin, S.Howison, M.Mineev, J.Ockendon, A.Orlov and M.Pavlov. The work of O.Y. and I.L. was supported by the European Community IIF's MIF1-CT-2004-007623 and MIF1-CT-2005-007323.

References

- [1] O. Agam, E. Bettelheim, P. Wiegmann, A. Zabrodin, Viscous fingering and a shape of an electronic droplet in the Quantum Hall regime, *Phys.Rev.Lett.* 88 (2002),cond-mat/0111333
- [2] G.Carlet, Extended Toda Hierarchy and its Hamiltonian Structure, PhD Thesis, SISSA preprint 2003/FM
- [3] Dickey, L. A. Soliton equations and Hamiltonian systems. Second edition. Advanced Series in Mathematical Physics, 26. World Scientific Publishing Co., Inc., River Edge, NJ, 2003
- [4] Dynamics of curved fronts. Edited by Pierre Pelc. Perspectives in Physics. Academic Press, Inc., Boston, MA, 1988. xvi+514 pp
- [5] B. Enriques , A. Orlov , V.Rubtsov , Dispersionful analogues of Benney's equations and N-wave systems ,*Inverse Problems* 12 (1996), no. 3, 241–250
- [6] Galin, L. A. Unsteady filtration with a free surface. *C. R. (Doklady) Acad. Sci. URSS (N.S.)* 47, (1945). 246–249
- [7] Howison, S. D. Fingering in Hele-Shaw cells. *J. Fluid Mech.* 167 (1986),
- [8] Kostov, I. K.; Krichever, I.; Mineev-Weinstein, M.; Wiegmann, P. B.; Zabrodin, A. The τ -function for analytic curves. *Random matrix models and their applications*, 285–299, *Math. Sci. Res. Inst. Publ.*, 40, Cambridge Univ. Press, Cambridge, 2001
- [9] I.Krichever, The τ -function of the universal Whitham hierarchy, matrix models and topological field theories. *Comm. Pure Appl. Math.* 47 (1994), no. 4, 437–475
- [10] M. Mineev-Weinstein, Selection of the Saffman-Taylor finger width in the absence of surface tension: an exact result, *Phys. Rev. Lett.* 80, 2113 (1998)
- [11] M. Mineev-Weinstein and S.P. Dawson, A new class of nonsingular exact solutions for Laplacian pattern formation, *Phys. Rev. E* 50, R24 (1994), *patt-sol/9305010* ;and *Phys. Rev. E* 57, 3063 (1998)
- [12] M. Mineev-Weinstein, P. Wiegmann, A. Zabrodin, Integrable Structure of Interface Dynamics, *Phys. Rev. Lett.* 84, (2000)5106-5109
- [13] Orlov, A. Yu.; Schulman, E. I. Additional symmetries for integrable equations and conformal algebra representation. *Lett. Math. Phys.* 12 (1986), no. 3, 171–179
- [14] Pavlov, M. V. Exact integrability of a system of Benney equations. (Russian) *Dokl. Akad. Nauk* 339 (1994), 311–313; translation in *Phys. Dokl.* 39 (1994), 745–747
- [15] Polubarinova-Kotschina, P. J. On the displacement of the oil-bearing contour. *C. R. (Doklady) Acad. Sci. URSS (N. S.)* 47, (1945). 250–254
- [16] Richardson S. Hele Shaw flows with a free boundary produced by the injection of fluid into a narrow channel. *J.Fluid Mech.*, 56, part 4, (1972), pp.609-618
- [17] Saffman, P. G.; Taylor, G., The penetration of a fluid into a porous medium or Hele-Shaw cell containing a more viscous liquid. *Proc. Roy. Soc. London. Ser., A* 245, 1958, 312–329. (2 plates)
- [18] K.Takasaki, T.Takebe, Integrable hierarchies and dispersionless limit. *Rev. Math. Phys.* 7 (1995), no. 5, 743–808

- [19] T.Takebe, Toda lattice hierarchy and conservation laws. Comm. Math. Phys. 129 (1990), no. 2, 281–318
- [20] Wiegmann, P. B.; Zabrodin, A., Conformal maps and integrable hierarchies, Comm. Math. Phys. 213 (2000), no. 3, 523–538